Generic Fincar (Finasteride): Uses, Benefits, Dosage, and Important Safety Information

Introduction

Generic Fincar (Finasteride) is a medication primarily used to manage symptoms associated with benign prostatic hyperplasia (BPH), a condition in which the prostate gland becomes enlarged. An enlarged prostate can make urination difficult and cause frequent trips to the bathroom, particularly during the night. Finasteride belongs to a group of medicines called 5-alpha-reductase inhibitors. It works by reducing the production of dihydrotestosterone (DHT), a hormone that contributes to prostate enlargement.

By lowering DHT levels, finasteride can gradually reduce prostate size, improve urinary flow, and help relieve symptoms associated with an enlarged prostate. Treatment generally requires consistent use over several months before noticeable improvements occur.

What Is Generic Fincar (Finasteride)?

Generic Fincar contains finasteride as its active ingredient. It is commonly available as a 5 mg oral tablet for treating benign prostatic hyperplasia. The medication works by blocking the activity of an enzyme called 5-alpha-reductase, which converts testosterone into DHT.

DHT plays an important role in prostate growth. When its production is reduced, the prostate may gradually become smaller. This can help reduce pressure on the urethra, the tube responsible for carrying urine out of the bladder.

Finasteride is also available in a 1 mg strength for male pattern hair loss, but the 5 mg strength is generally used for prostate-related conditions. These strengths are intended for different treatment purposes and should not be interchanged without medical advice.

Uses of Generic Fincar

The primary use of finasteride 5 mg is the treatment of benign prostatic hyperplasia in adult men. Its potential benefits include:

- **Reducing prostate size:** Helps gradually shrink an enlarged prostate by lowering DHT levels.

- **Improving urine flow:** May make urination easier by reducing pressure on the urethra.
- **Managing urinary symptoms:** Can help reduce difficulty starting urination, weak urine flow, and incomplete bladder emptying.
- **Reducing urinary retention risk:** Long-term treatment can lower the likelihood of acute urinary retention.
- **Reducing the need for surgery:** In suitable patients, treatment may decrease the likelihood of requiring prostate surgery.

How Does Finasteride Work?

Finasteride works by inhibiting the type II 5-alpha-reductase enzyme. This enzyme is responsible for converting testosterone into dihydrotestosterone (DHT). Higher DHT activity can contribute to prostate tissue growth.

When finasteride reduces DHT production, the prostate may gradually decrease in size. As the pressure around the urethra decreases, urine can pass more easily. Unlike some medicines that provide relatively quick relief by relaxing prostate muscles, finasteride works gradually by addressing hormonal activity.

FDA Access Data+1

Recommended Dosage and Administration

For benign prostatic hyperplasia, the commonly recommended finasteride dosage is 5 mg once daily. It can generally be taken with or without food. The tablet should be swallowed with water and taken according to the healthcare professional's instructions.

Patients should maintain a consistent medication schedule and avoid taking extra tablets if a dose is missed. The medicine may require three to six months or longer to provide noticeable benefits.

The prescribed dosage and treatment duration should always be determined by a qualified healthcare professional.

DailyMed+1

Possible Side Effects

Like other prescription medicines, finasteride may cause side effects in some individuals. Not everyone experiences these reactions, and their intensity can vary.

Some reported side effects include:

- Reduced sexual desire or libido.

- Difficulty achieving or maintaining an erection.
- Changes in ejaculation or reduced semen volume.
- Breast tenderness or enlargement.
- Skin rash or allergic reactions.

Patients should inform their healthcare professional about persistent or concerning symptoms. Breast lumps, nipple discharge, or unusual breast changes should receive medical evaluation.

DailyMed+1

Precautions and Safety Information

Finasteride should be used under medical supervision. Patients with liver problems, urinary difficulties, or a history of allergic reactions to the medication should discuss these conditions before starting treatment.

Pregnant women or women who may become pregnant should not handle crushed or broken finasteride tablets because exposure can pose a risk to the development of a male fetus. The medication is not indicated for women or children.

Finasteride can also reduce prostate-specific antigen (PSA) levels, which are measured during prostate health assessments. Therefore, patients should inform their healthcare professional that they are taking this medicine before undergoing PSA testing. Regular medical follow-ups help monitor treatment progress and prostate health.

FINASTERIDE tablet, film coated+1

Storage Instructions

Store finasteride tablets at room temperature in a dry place, away from excessive heat, moisture, and direct sunlight. Keep the medicine in its original packaging and out of reach of children. Always check the product packaging for specific storage recommendations and expiry information.

Conclusion

Generic Fincar (Finasteride) is a prescription medication primarily used to manage symptoms of benign prostatic hyperplasia. By reducing DHT production, it helps gradually decrease prostate size, improve urinary flow, and lower the risk of certain complications associated with prostate enlargement. Consistent use, medical supervision, and regular follow-ups are important for safe and effective treatment. Patients should understand the possible side effects and follow their prescribed dosage rather than adjusting treatment independently.